

Dysport.
(abobotulinumtoxinA)

Uniquely You. Definitely Dysport®.

Get the injection for frown lines between the brows that delivers natural-looking, long-lasting results visible as early as 1 day after treatment.^{1-5*}



Actual patient. Individual results may vary.

Dysport® (abobotulinumtoxinA) is a prescription injection for temporary improvement in the look of moderate to severe frown lines between the eyebrows (glabellar lines) in adults less than 65 years of age.

Please see full Important Safety Information, including Distant Spread of Toxin Effect Boxed Warning, on pages 6 and 7.

*GL-1 and GL-3 evaluated subjects for at least 150 days following treatment. Based on a ≥1-Grade GLSS improvement from baseline utilizing data from 2 double-blinded, randomized, placebo-controlled pivotal studies (GL-1, GL-3) in a post hoc analysis.

Natural-looking. Fast-acting. Long-lasting.^{2-5*}

With more than 30 years of clinical experience, an established safety profile, and over 100 million treatments worldwide, *Dysport* delivers results that look and feel like you.^{6,7}

98%
of users

98% of users say they would get *Dysport* again, while 97% of users say their results look natural.^{8†}

1
day

You may see results as early as 1 day after treatment, with typical results within 2 to 3 days.^{2-4‡}

5
months

Improvements to frown lines can last up to 5 months.^{2-5*†}

Definitely satisfied.

95% of users were satisfied with only 2 treatments per year.^{8†}

Please see full Important Safety Information, including Distant Spread of Toxin Effect Boxed Warning, on pages 6 and 7.

*GL-1 and GL-3 evaluated subjects for at least 150 days following treatment. Based on a ≥1-Grade GLSS improvement from baseline utilizing data from 2 double-blinded, randomized, placebo-controlled pivotal studies (GL-1, GL-3) in a post hoc analysis.

†Users = clinical trial subjects. Subject-reported at 12 months (N=120) after 2 treatments 6 months apart in a phase IV, multicenter, prospective study.

‡Based on subject self-assessment. The onset of response at day 1 was 15% (16/105) in GL-1 and 33% (65/200) in GL-3. In the DREAM study, the onset of response at day 1 was 33% (66/200). The median time to onset of response was 3 days in GL-1 (55/105; 52%) and GL-2 (36/71; 51%), and 2 days in GL-3 (110/200; 55%).

§Subject-reported at 12 months (N=120) after 2 treatments 6 months apart in a phase IV, multicenter, prospective study.

Smooth between-the-brow frown lines

Dysport blocks the signal from the nerve to the muscles, which temporarily prevents contractions of the muscles that cause frown lines.⁹

Untreated muscles still work normally, so you can continue to express yourself naturally and confidently.⁵




Actual patient.
Individual results may vary.

Dysport.
(abobotulinumtoxinA)

Enhance your look and boost your confidence

Dysport produces similar results with repeat treatments, giving you a natural-looking confidence boost again and again.^{2,10,11†}

Treatment area



One injection into each of the 5 points between and above the eyebrows temporarily prevents muscle contractions.¹

Rebecca's look with Dysport

Before

After



Frown after 50 units, day 27

Please see full Important Safety Information, including Distant Spread of Toxin Effect Boxed Warning, on pages 6 and 7.

[†]Subject-reported at 12 months (N=120) after 2 treatments 6 months apart in a phase IV, multicenter, prospective study.



Actual patient.
Individual results may vary.

Dysport.
(abobotulinumtoxinA)

Brief Summary

This summary contains important information about *Dysport* (DIS–port) for injection. Read this information carefully before using *Dysport*. Ask your doctor if you do not understand any of this information or if you want to know more about *Dysport*. For full Prescribing Information and Medication Guide, please see the package insert.

What is *Dysport*?

Dysport is a prescription injection for temporary improvement in the look of moderate to severe frown lines between the eyebrows (glabellar lines) in adults less than 65 years of age.

Important Safety Information

WARNING: DISTANT SPREAD OF TOXIN EFFECT

In some cases, the effects of *Dysport* and all botulinum toxin products may affect areas of the body away from the injection site. Symptoms can happen hours to weeks after injection and may include swallowing and breathing problems, loss of strength and muscle weakness all over the body, double vision, blurred vision and drooping eyelids, hoarseness or change or loss of voice, trouble saying words clearly, or loss of bladder control. Swallowing and breathing problems can be life threatening and there have been reports of death. You are at the highest risk if these problems are pre-existing before injection. These effects could make it unsafe for you to drive a car, operate machinery, or do other dangerous activities.

Who should not take *Dysport*?

Do not take *Dysport* if you:

- are allergic to *Dysport* or any of its ingredients. See the end of this Medication Guide for a list of ingredients in *Dysport*
- are allergic to cow's milk protein
- had an allergic reaction to any other botulinum toxin product such as Myobloc® (rimabotulinumtoxinB), Botox® (onabotulinumtoxinA), or Xeomin® (incobotulinumtoxinA)
- have a skin infection at the planned injection site
- under 18 years of age
- pregnant or breastfeeding

What additional warnings and precautions should I know about *Dysport*?

- **The dose of *Dysport* is not the same as the dose of any other botulinum toxin product and cannot be compared to the dose of any other botulinum toxin product you may have used.**
- *Dysport* may not be right for you if you have surgical changes to your face, very weak muscles in the treatment area, your face looks very different from side to side, the injection site is inflamed, you have droopy eyelids or sagging eyelid folds, deep facial scars, thick oily skin, or if your wrinkles can't be smoothed by spreading them apart.
- **Tell your doctor about all of your medical conditions**, especially if you have a disease that affects muscles and nerves (such as amyotrophic lateral sclerosis [ALS or Lou Gehrig's disease], myasthenia gravis, or Lambert-Eaton syndrome), allergies to any botulinum toxin product or had any side effect from any botulinum toxin product in the past, a breathing problem (such as asthma or emphysema), swallowing problems, bleeding problems, diabetes, or a slow heart beat or other problem with your heart rate or rhythm, plans to have surgery, had surgery on your face, have weakness in or near your muscles being treated, drooping eyelids, any other change in the way your face normally looks, or are pregnant or breastfeeding or planning to become pregnant or breastfeed. Patients with swallowing or breathing difficulties or a disease that affects muscles and nerves may have a higher risk of serious side effects, including severe swallowing and breathing problems, after treatment with *Dysport*.
- **Tell your doctor about all the medicines you take**, including prescription and nonprescription medicines, vitamins and herbal and other natural products. Using *Dysport* with certain other medicines may cause serious side effects. **Do not start any new medicines while taking *Dysport* without talking to your doctor first.**
- **Especially tell your doctor if you:** have received any other botulinum toxin product, such as Myobloc® (rimabotulinumtoxinB), Botox® (onabotulinumtoxinA), or Xeomin® (incobotulinumtoxinA), in the last

four months or any in the past (be sure your doctor knows exactly which product you received, have recently received an antibiotic by injection, take muscle relaxants, take an allergy or cold medicine, or take a sleep medicine.

What are the side effects of *Dysport*?

The most common side effects of *Dysport* include nose and throat irritation, injection site pain, upper respiratory infection, blood in urine, headache, injection site reaction, eyelid swelling, eyelid drooping, sinus infection, and nausea.

These are not all the possible side effects of *Dysport*. Call your doctor for medical advice about side effects.

You may report side effects to FDA at www.fda.gov/medwatch or 1-800-FDA-1088.

This Brief Summary summarizes the most important information about *Dysport*. If you would like more information, talk with your doctor.

What are the ingredients of *Dysport*?

Active ingredient: (botulinum toxin Type A)

Inactive ingredients: human albumin and lactose. *Dysport* may contain cow's milk protein.

Where should I go for more information about *Dysport*?

- Talk to your doctor.
- Visit www.DYSPORTUSA.com.

GALDERMA LABORATORIES, L.P.

Dallas, Texas 75201 USA

Revised: September 2023


References

1. *Dysport* prescribing information. Galderma Laboratories, LP; 2023.
2. Rubin MG, Dover J, Glogau RG, Goldberg DJ, Goldman MP, Schlessinger J. The efficacy and safety of a new U.S. Botulinum toxin type A in the retreatment of glabellar lines following open-label treatment. *J Drugs Dermatol*. 2009;8(5):439–444.
3. Monheit GD, Baumann L, Maas C, Rand R, Down R. Efficacy, safety, and subject satisfaction after abobotulinumtoxinA treatment for moderate to severe glabellar lines. *Dermatol Surg*. 2020;46(1):61–69.
4. Brandt F, Swanson N, Baumann L, Huber B. Randomized, placebo-controlled study of a new botulinum toxin type A for treatment of glabellar lines: efficacy and safety. *Dermatol Surg*. 2009;35(12):1893–1901.
5. Data on file. MA-35497. Post hoc analysis. Fort Worth, TX: Galderma Laboratories, L.P., 2017.
6. Electronic Medicines Compendium. *Dysport* 500 units powder for solution for injection. Updated August 1, 2024. Accessed November 24, 2024. <https://www.medicines.org.uk/emc/product/7261/smpc>
7. Data on file. MA-52573. More than 100 million *Dysport* Azzalure treatments delivered in aesthetics indications.
8. Schlessinger J, Cohen JL, Shamban A, et al. A multicenter study to evaluate subject satisfaction with two treatments of AbobotulinumtoxinA a year in the glabellar lines. *Dermatol Surg*. 2021;47(4):504–509.
9. Molina B, Grangier Y, Mole B, et al. Patient satisfaction after the treatment of glabellar lines with Botulinum toxin type A (Speywood Unit): a multi-centre European observational study. *J Eur Acad Dermatol Venereol*. 2015;29(7):1382–1388.
10. Moy R, Maas C, Monheit G, Huber MB; Reloxin Investigational Group. Long-term safety and efficacy of a new botulinum toxin type A in treating glabellar lines. *Arch Facial Plast Surg*. 2009;11(2):77–83.
11. Cohen JL, Schlessinger J, Cox SE, Lin X; Reloxin Investigational Group. An analysis of the long-term safety data of repeat administrations of botulinum neurotoxin type A-ABO for the treatment of glabellar lines. *Aesthet Surg J*. 2009;29(6 Suppl):S43–S49.
12. Kane MA, Monheit G. The practical use of abobotulinumtoxinA in aesthetics. *Aesthet Surg J*. 2017;37(suppl 1):S12–S19.
13. Small R. Botulinum toxin injection for facial wrinkles. *Am Fam Physician*. 2014;90(3):168–175.
14. King M. The management of bruising following nonsurgical cosmetic treatment. *J Clin Aesthet Dermatol*. 2017;10(2):E1–E4.

Dysport.
(abobotulinumtoxinA)


How to get the best out of your *Dysport* treatment

Proper aftercare helps prevent unwanted spread and lowers the risk of bruising.



For the first 4 hours after treatment, avoid:

- Lying flat^{12,13}
- Rubbing or massaging injected areas^{12,13}



For the first 24 hours after treatment, avoid:

- Intense exercise or activities that increase blood flow^{12,14}
- Exposure to hot temperatures or activities that cause flushing^{13,14}

ASPIRE

GALDERMA REWARDS



See the results.
Get the rewards.

Sign up for ASPIRE Galderma Rewards to earn points on qualifying *Dysport* treatments and save on future services in your aesthetic journey, including Restylane® and Sculptra®.

Dysport.
(abobotulinumtoxinA)

GALDERMA
EST. 1981

Please see full Important Safety Information, including Distant Spread of Toxin Effect Boxed Warning, on pages 6 and 7.

© 2025 Galderma Laboratories, L.P. The *Dysport* trademark is used under license. All trademarks are the property of their respective owners. US-DYS-2300161